

Constraints on the gluon Siverts distribution via transverse single spin asymmetries at midrapidity in $p^\uparrow p \rightarrow \pi^0 X$ processes at BNL RHIC

M. Anselmino,^{1,*} U. D'Alesio,^{2,†} S. Melis,^{2,‡} and F. Murgia^{2,§}

¹*Dipartimento di Fisica Teorica, Università di Torino and Istituto Nazionale di Fisica Nucleare, Sezione di Torino, Via P. Giuria 1, I-10125 Torino, Italy*

²*Istituto Nazionale di Fisica Nucleare, Sezione di Cagliari and Dipartimento di Fisica, Università di Cagliari, C.P. 170, I-09042 Monserrato (CA), Italy*

(Dated: February 2, 2008)

We consider the recent RHIC data on the transverse single spin asymmetry (SSA) A_N , measured in $p^\uparrow p \rightarrow \pi^0 X$ processes at mid-rapidity by the PHENIX collaboration. The measurement is consistent with a vanishing SSA. We analyze this experimental information within a hard scattering approach based on a generalized QCD factorization scheme, with unintegrated, transverse momentum dependent (TMD), parton distribution and fragmentation functions. It turns out that, in the kinematical region of the data, only the gluon Siverts effect could give a large contribution to A_N ; its vanishing value is thus an indication about the possible size of the gluon Siverts function (GSF). Approximate upper limits on its magnitude are derived. Additional constraints obtained combining available parameterizations of the quark Siverts function and the Burkardt sum rule (BSR) for the Siverts distributions are also discussed.

PACS numbers: 12.38.Bx, 13.88.+e, 13.85.Ni, 14.70.Dj

I. INTRODUCTION AND APPROACH

We have recently discussed a hard scattering approach to hadronic interactions, based on the assumption of a generalized QCD factorization scheme which involves unintegrated TMD parton distribution and fragmentation functions; the partonic intrinsic motions are also fully taken into account in the elementary perturbative QCD (pQCD) processes [1, 2, 3]. This scheme has been applied to the computation of unpolarized cross sections for single-inclusive particle production, $pp \rightarrow h X$, at high energy and (moderately) large p_T [1], and to the computation of the transverse SSA

$$A_N = \frac{d\sigma^\uparrow - d\sigma^\downarrow}{d\sigma^\uparrow + d\sigma^\downarrow} \quad (1)$$

in $p^\uparrow p \rightarrow h X$ processes [1, 2, 3, 4].

Transverse single spin asymmetries can originate, even with a short distance helicity conserving pQCD dynamics, from spin- $\mathbf{k}_\perp$ correlations in the soft components of the hadronic process, the distribution and fragmentation functions. There could be many such correlations. However, the study of SSA based on the assumption of a generalized factorization scheme [2, 3] shows that the correct treatment of the elementary pQCD dynamics, with all phases related to the non-collinear and non-planar partonic configurations, leads to a strong suppression of all contributions, with the exception of the Siverts [5], and, to a lesser extent, the Collins [6] mechanisms.

The issue of the validity of the factorization scheme with unintegrated partonic distribution and fragmentation functions is still an open one. Such a scheme has been shown to hold for Semi-Inclusive Deep Inelastic Scattering (SIDIS) and Drell-Yan processes [7, 8, 9], while it is not yet clear whether or not it holds for inclusive one-particle production in pp processes; in such a case it is difficult to account for the gauge links necessary to ensure the gauge invariance of the TMD distribution functions [10, 11, 12, 13, 14]. At this stage we consider our factorized description of $pp \rightarrow \pi X$ processes as a phenomenological model based on a natural extension of the usual collinear approach for the same process, and of the factorized scheme with unintegrated partonic distributions proven for SIDIS and Drell-Yan processes.

The Siverts function has recently received a lot of attention: data on azimuthal SSA in semi-inclusive deep inelastic scattering (SIDIS) processes from the HERMES collaboration at DESY [15] and from the COMPASS collaboration

*Electronic address: mauro.anselmino@to.infn.it

†Electronic address: umberto.dalesio@ca.infn.it

‡Electronic address: stefano.melis@ca.infn.it

§Electronic address: francesco.murgia@ca.infn.it

at CERN [16] have allowed, for the first time, a direct extraction of the Sivvers functions for u and d quarks inside a proton [17, 18, 19, 20, 21]. Similarly, u and d Sivvers functions have been extracted from purely hadronic processes [1], in qualitative agreement with the SIDIS results. Also the possibility of accessing the gluon Sivvers function has been investigated [4, 22, 23, 24]. Although one might expect that spin- $\mathbf{k}_\perp$ correlations are stronger for valence quarks – as the large x_F data from the FNAL-E704 collaboration [25] and the STAR collaboration at RHIC-BNL [26] seem to indicate – one knows that gluons play a dominant role in many high energy hadronic processes; it would be very interesting to see whether or not the gluon density inside a transversely polarized proton depends on the intrinsic motion.

We address here the issue of the largely unknown gluon Sivvers function, $\Delta^N \hat{f}_{g/p^\uparrow}(x, \mathbf{k}_\perp)$, and its possible contribution to the SSA A_N for the $p^\uparrow p \rightarrow \pi^0 X$ process, in the framework of the generalized factorization scheme, with pQCD elementary dynamics, of Refs. [1, 2, 3]. In this approach, the general structure of the cross section for the polarized hadronic process $(A, S_A) + (B, S_B) \rightarrow C + X$, is given by [3]

$$\begin{aligned} \frac{E_C d\sigma^{(A,S_A)+(B,S_B) \rightarrow C+X}}{d^3\mathbf{p}_C} &= \sum_{a,b,c,d,\{\lambda\}} \int \frac{dx_a dx_b dz}{16\pi^2 x_a x_b z^2 s} d^2\mathbf{k}_{\perp a} d^2\mathbf{k}_{\perp b} d^3\mathbf{k}_{\perp C} \delta(\mathbf{k}_{\perp C} \cdot \hat{\mathbf{p}}_c) J(k_{\perp C}) \\ &\times \rho_{\lambda_a, \lambda'_a}^{a/A, S_A} \hat{f}_{a/A, S_A}(x_a, \mathbf{k}_{\perp a}) \rho_{\lambda_b, \lambda'_b}^{b/B, S_B} \hat{f}_{b/B, S_B}(x_b, \mathbf{k}_{\perp b}) \\ &\times \hat{M}_{\lambda_c, \lambda_d; \lambda_a, \lambda_b}^* \hat{M}_{\lambda'_c, \lambda'_d; \lambda'_a, \lambda'_b} \delta(\hat{s} + \hat{t} + \hat{u}) \hat{D}_{\lambda_c, \lambda'_c}^{\lambda_c, \lambda'_c}(z, \mathbf{k}_{\perp C}), \end{aligned} \quad (2)$$

where all parton intrinsic motions are fully taken into account, both in the soft, non perturbative components and in the hard, pQCD interactions.

In Ref. [3] the structure of Eq. (2) was extensively discussed; its main features are the appearance of several spin and $\mathbf{k}_\perp$ dependent distribution and fragmentation functions (with a partonic interpretation) and the non-collinear partonic configurations which lead to many $\mathbf{k}_\perp$ dependent phases. As a consequence, it was explicitly shown how the integration over the parton intrinsic momenta leads to strong suppressions of most contributions to the unpolarized cross sections and to the transverse single spin asymmetry A_N . The only sizeable contributions to the latter, in the kinematical region (large positive x_F) of the E704 [25] and STAR data [26] come from the Sivvers and, less importantly, from the Collins mechanisms. The dominance of the Sivvers effect is even more pronounced in other x_F ranges, namely at $x_F \leq 0$.

In Ref. [3] also the flavour decomposition of the Sivvers effect was performed; while the quark contribution is totally dominant at large and positive x_F values (for polarized protons moving along the positive Z -axis), the gluon contribution may be sizeable at $x_F \simeq 0$, the mid-rapidity region. This can be easily understood, since $x_a^{\min}$ (the lowest kinematically accessible value of the light-cone momentum fraction of parton a inside the transversely polarized initial proton) must be larger than x_F ; A_N at large x_F is then mainly driven by valence quark properties. Indeed, an analysis of the E704 data allowed a first extraction of the Sivvers functions for u and d quarks [1]. In principle, also inclusive production in the negative x_F region might be sensitive to small x_a gluons inside the polarized proton (being hit by large x_b partons inside the unpolarized one). We shall further comment on this point after Eq. (5).

Data in the mid-rapidity region are available from the E704 [27] and PHENIX [28] experiments. The kinematical regime corresponding to negative values of x_F has been covered by the STAR collaboration [29]. Preliminary results for charged pions are also available from the RHIC-BRAHMS experiment [30].

In these kinematical regions $A_N(p^\uparrow p \rightarrow \pi X)$ is largely dominated by the Sivvers effect alone, and Eq. (2) gives [1]:

$$\begin{aligned} \frac{E_\pi d\sigma^\uparrow}{d^3\mathbf{p}_\pi} - \frac{E_\pi d\sigma^\downarrow}{d^3\mathbf{p}_\pi} &\simeq \sum_{a,b,c,d} \int \frac{dx_a dx_b dz}{\pi x_a x_b z^2 s} d^2\mathbf{k}_{\perp a} d^2\mathbf{k}_{\perp b} d^3\mathbf{k}_{\perp \pi} \delta(\mathbf{k}_{\perp \pi} \cdot \hat{\mathbf{p}}_c) J(k_{\perp \pi}) \\ &\times \Delta \hat{f}_{a/p^\uparrow}(x_a, \mathbf{k}_{\perp a}) \hat{f}_{b/p}(x_b, \mathbf{k}_{\perp b}) \hat{s}^2 \frac{d\hat{\sigma}^{ab \rightarrow cd}}{d\hat{t}}(x_a, x_b, \hat{s}, \hat{t}, \hat{u}) \delta(\hat{s} + \hat{t} + \hat{u}) \hat{D}_{\pi/c}(z, \mathbf{k}_{\perp \pi}), \end{aligned} \quad (3)$$

where (M denotes the proton mass)

$$\Delta \hat{f}_{a/p^\uparrow}(x_a, \mathbf{k}_{\perp a}) \equiv \hat{f}_{a/p^\uparrow}(x_a, \mathbf{k}_{\perp a}) - \hat{f}_{a/p^\downarrow}(x_a, \mathbf{k}_{\perp a}) = \Delta^N \hat{f}_{a/p^\uparrow}(x_a, k_{\perp a}) \cos \phi_a = -2 \frac{k_{\perp a}}{M} f_{1T}^\perp(x_a, k_{\perp a}) \cos \phi_a. \quad (4)$$

$\Delta^N \hat{f}_{a/p^\uparrow}(x_a, k_{\perp a})$ [or $f_{1T}^\perp(x_a, k_{\perp a})$] is referred to as the Sivvers distribution function of parton a inside a transversely polarized proton. ϕ_a is the azimuthal angle of the intrinsic transverse momentum $\mathbf{k}_{\perp a}$ of parton a . We follow here the notations and kinematical conventions of Ref. [3], with the polarized proton moving along the positive Z -axis, in the pp c.m. frame; the observed pion is produced in the XZ plane, with positive X values; spin $\uparrow$ and $\downarrow$ are respectively along the positive and negative Y -axis.

Eq. (3) gives the numerator of A_N , Eq. (1); the denominator is just twice the unpolarized cross section, which is given by

$$\begin{aligned} \frac{E_\pi d\sigma^{unp}}{d^3\mathbf{p}_\pi} &\simeq \sum_{a,b,c,d} \int \frac{dx_a dx_b dz}{\pi x_a x_b z^2 s} d^2\mathbf{k}_{\perp a} d^2\mathbf{k}_{\perp b} d^3\mathbf{k}_{\perp\pi} \delta(\mathbf{k}_{\perp\pi} \cdot \hat{\mathbf{p}}_c) J(k_{\perp\pi}) \\ &\times \hat{f}_{a/p}(x_a, k_{\perp a}) \hat{f}_{b/p}(x_b, k_{\perp b}) \hat{s}^2 \frac{d\hat{\sigma}^{ab \rightarrow cd}}{d\hat{t}}(x_a, x_b, \hat{s}, \hat{t}, \hat{u}) \delta(\hat{s} + \hat{t} + \hat{u}) \hat{D}_{\pi/c}(z, k_{\perp\pi}). \end{aligned} \quad (5)$$

The azimuthal phase factor $\cos\phi_a$ appearing in the numerator of A_N , Eqs. (3) and (4), plays a crucial role and deserves a comment. The only other term depending on ϕ_a in Eq. (3) is the partonic cross section, in particular via the corresponding Mandelstam variable $\hat{t}$. At large positive x_F and moderately large p_T the (average) values of $\hat{t}$ are relatively small. Therefore, the (dominant) $\hat{t}$ -channel contributions, proportional to $1/\hat{t}^2$, depend sizeably on ϕ_a , so that the $d^2\mathbf{k}_{\perp a} \cos\phi_a/\hat{t}^2$ integration in Eq. (3) does not necessarily suppress A_N . Instead, for negative values of x_F , all partonic Mandelstam variables are much less dependent on ϕ_a , so that one is roughly left with the $d^2\mathbf{k}_{\perp a} \cos\phi_a$ integration alone, which cancels the potentially large Siverts contribution. As a consequence, the possibility of gaining information on the gluon Siverts distribution from the recent STAR and BRAHMS data at negative values of x_F is frustrated. Notice that, due to the much lower values of $\sqrt{s}$ involved, this suppression caused by the $\cos\phi_a$ dependence would be much less effective for the kinematical regimes of E704 and of the proposed PAX [31] experiments [3]. However, E704 never measured A_N for negative x_F values, while the PAX experiment is still in the proposal and planning stage.

Similar considerations lead to a strong suppression of the gluon Siverts contribution to the SSA for $p^\uparrow p \rightarrow \gamma X$ processes at STAR, in the negative x_F range and at a (pseudo)rapidity $\eta \simeq -4$ [32].

The same arguments do not apply to inclusive hadronic processes at mid-rapidity and moderately large p_T values, for which data from PHENIX [28] are already available, for neutral pions and charged hadron production. For these processes, the gluon contribution is dominant and the Siverts effect can survive the phase integration. In the next section, we shall therefore consider in detail this case, aiming at a possible derivation, from data, of useful direct constraints on the gluon Siverts function.

A. Constraints from the Burkardt sum rule

Indirect constraints on the GSF could also be obtained from a sum rule for the Siverts distribution recently derived by Burkardt [33]. The BSR states that the total (integrated over x and $\mathbf{k}_\perp$) transverse momentum of all partons (quarks, antiquarks and gluons) in a transversely polarized proton must be zero,

$$\langle \mathbf{k}_\perp \rangle = \sum_a \langle \mathbf{k}_\perp \rangle_a = \int dx \int d^2\mathbf{k}_\perp \mathbf{k}_\perp \sum_a \Delta \hat{f}_{a/p^\uparrow}(x, \mathbf{k}_\perp) = 0. \quad (6)$$

Naively, at the partonic level considered in our approach, this sum rule simply corresponds to total (transverse) momentum conservation inside a transversely polarized proton, since the unintegrated distribution function

$$\hat{f}_{a/p^\uparrow}(x_a, \mathbf{k}_{\perp a}) = \hat{f}_{a/p}(x_a, k_{\perp a}) + \frac{1}{2} \Delta \hat{f}_{a/p^\uparrow}(x_a, \mathbf{k}_{\perp a}), \quad (7)$$

introduced in Eq. (4), has a clear probabilistic interpretation. On the other hand, it is well known [10, 11] that a non-vanishing Siverts effect requires initial/final state interactions, which might spoil the simple partonic interpretation. The Burkardt sum rule ensures the non-trivial result that momentum conservation holds also in this situation. The validity of the BSR has been explicitly verified in a diquark spectator model calculation of the Siverts distribution in Ref. [34]. In this paper, we consider the BSR as an additional theoretical tool in order to obtain indirect information on the gluon Siverts distribution, once the quark distributions are known.

We notice, however, that a strict use of the BSR requires integration over the full x range, including the poorly known small x region, of the single Siverts functions, which might even result in divergences. Therefore, in the following we shall simply check whether or not the parameterizations of the Siverts functions, which will turn out from our phenomenological analysis of the data within the theoretical approach of Refs. [1, 2, 3], fulfill the BSR in the limited x range considered, without extrapolating them to very low x values.

In the next section we present our results on the GSF, based on the PHENIX [28] experimental data on $A_N(p^\uparrow p \rightarrow \pi^0 X)$. Further comments and conclusions are given in the last section.

II. PHENOMENOLOGY OF THE GLUON SIVERS FUNCTION

We consider the PHENIX data [28] on A_N for the $p^\uparrow p \rightarrow \pi^0 X$ process at RHIC, at $\sqrt{s} = 200$ GeV, with p_T ranging from 1.0 to 5.0 GeV/ c and mid-rapidity values, $|\eta| < 0.35$. In this kinematical regime, at the lowest p_T values, $x_a^{\min}$ can be as small as 0.005. Therefore, partonic channels involving a gluon in the transversely polarized initial proton dominate over those involving a quark. This gluon dominance, together with the (almost) vanishing of all possible contributions to A_N other than the Sivers effect, allow to interpret the data – showing tiny values of A_N – in terms of useful constraints (upper bounds) on the gluon Sivers function, as it will be shown. As p_T grows, $x_a^{\min}$ increases and the dominance of the gluon channels becomes less prominent.

Another set of data, for comparable rapidity and p_T ranges, has been collected several years ago by the E704 Collaboration [27], at a lower energy, $\sqrt{s} \simeq 20$ GeV. In this case, however, even at the smallest p_T values, $x_a^{\min}$ remains large enough so that gluon channels are not dominating; a possible mixing with quark initiated contributions then prevents to get clean constraints on the GSF from E704 lower energy data.

Before analyzing in detail the reliable (for our purpose) PHENIX data, let us summarize what we know so far about the Sivers functions. As already discussed, the E704 and STAR collaborations have measured large SSA for the $p^\uparrow p \rightarrow \pi X$ process at large positive x_F and moderate p_T values. In this kinematical regime, since $x_a^{\min} > x_F$, gluon and sea-quark contributions should be negligible. Indeed, in Ref. [1] we have shown that reasonable fits to the SSA can be obtained by using valence-like Sivers functions for u and d quarks, which turn out to have opposite signs. In addition, fits to the weighted azimuthal asymmetries measured for pion production in SIDIS with a transversely polarized target [15, 16], lead to independent comparable parameterizations of the u and d Sivers distributions [17, 18, 20, 21].

Notice that in all these phenomenological analyses the Sivers functions for gluons and sea-quarks have been assumed to be vanishing. That is a reasonable assumption, considering the kinematical region of the available data, and the lowest order decoupling of gluons in SIDIS. A natural question arising at this point is the following: how do the valence u and d Sivers functions alone, so far extracted, perform with the mid-rapidity PHENIX and E704 data? The answer is that they predict an almost vanishing SSA, compatible with both sets of data (see Fig. 1, for the PHENIX results). Not only, but their parameterizations [1, 17] are also compatible with the Burkardt sum rule, Eq. (6), yielding $\langle \mathbf{k}_\perp \rangle \simeq 0$ within a 10% accuracy. It looks like there is no need to introduce other contributions to the Sivers effect in addition to that coming from valence quarks.

Of course, this cannot be a definite, although simple and appealing, conclusion, as the data used for the extraction of the u and d quark Sivers functions are insensitive to the small x region. A large gluon Sivers function would not affect the analysis of the SIDIS, E704 and STAR data at large positive x_F ; however, it would strongly affect the description of the mid-rapidity PHENIX data. The real question is now: how much does the small value of A_N measured by PHENIX suppress the gluon Sivers function?

In order to answer this question we compute A_N according to Eqs. (1) and (3)-(5), imposing different conditions on the gluon Sivers functions, trying to understand what is the maximum value of $|\Delta^N \hat{f}_{g/p^\uparrow}(x, k_\perp)|/2\hat{f}_{g/p}(x, k_\perp)$ allowed by the PHENIX data. We follow Ref. [1], adopting the same factorized gaussian $k_\perp$ dependence for distribution and fragmentation functions and the same parameterization for the Sivers distributions, which are related to the unpolarized parton densities; the latter are given by the MRST01 set [35], and the fragmentation functions by the KKP set [36]. The valence u and d quark Sivers functions are the same as extracted from E704 data in Ref. [1].

Our results are shown in Figs. 1 and 2 and require some comments.

- The thin, red, solid line in Fig. 1, is the result of computing A_N using only the valence u and d Sivers functions, as extracted in Ref. [1].
- The cyan, dot-dashed curve in Fig. 1 corresponds to the extreme case of the largest (in magnitude) gluon Sivers distribution, obtained by saturating the natural positivity bound, see Eq. (4),

$$\Delta^N \hat{f}_{g/p^\uparrow}(x, k_\perp) = -2\hat{f}_{g/p}(x, k_\perp). \quad (8)$$

The sea-quark Sivers functions are assumed to vanish. This leads to a SSA definitely in contradiction with the data. It also leads to a strong violation of the BSR. As expected, from PHENIX data a much more stringent constraint than the simple positivity bound for the (absolute value of the) gluon Sivers function can be obtained. The maximized (in magnitude) GSF in Eq. (8) has been chosen negative as the data hint at a possible small negative value of A_N . The opposite choice would lead to very similar conclusions, actually with an even stronger disagreement with data.

- The thick, red, solid curve in Fig. 1 has been obtained still assuming that there is no sea-quark Sivers contribution, and looking for a parameterization of $\Delta^N \hat{f}_{g/p^\uparrow}$ yielding values of A_N falling, approximately, within one-sigma deviation below the lowest p_T data. This somehow corresponds to the largest (in magnitude) acceptable gluon Sivers function, in the sense that any larger GSF would cause the SSA to lie outside the error bars.

The corresponding x -dependent part of the GSF, normalized to its positivity bound, $|\Delta^N f_{g/p^\uparrow}(x)|/2f_{g/p}(x)$, is shown as the red, solid curve in Fig. 2, as a function of x . PHENIX data on A_N clearly impose a stringent upper

bound on the magnitude of the gluon Sivvers function in the region of small x , where gluons play a crucial role. The constraint is much less significant at larger x values, as it is natural: there cannot be any strong correlation between gluon distributions at large x (where they are anyway negligible), and a physical observable, like A_N at mid-rapidity, which is mainly sensitive to small x values.

The GSF obtained here, the red, solid curve of Fig. 2, leads, within the x range covered by the data, to a strong violation of the BSR. This could be avoided by imposing an even smaller (in size) GSF, giving a strong role to the BSR.

- As a last attempt, we have released the assumption of vanishing sea-quark Sivvers distributions. With the aim of exploring how large a GSF can be, we have done that in an extreme scenario, *i.e.* assuming that all sea-quarks (u_s , $\bar{u}$, d_s , $\bar{d}$, s , $\bar{s}$) have a non-vanishing positive Sivvers function which saturates the positivity bound [that is, $\Delta^N \hat{f}_{q_s/p^\uparrow}(x, k_\perp) \equiv 2\hat{f}_{q_s/p}(x, k_\perp)$]. Their contribution could then cancel the negative contribution to A_N of a possibly large GSF.

Again, we look for the largest negative GSF which, together with a positive maximized sea-quark contribution, gives a SSA approximately lying within one-sigma standard deviation below the data points, represented again by the thick, red, solid line of Fig. 1. This curve results now as the sum of the (maximized) sea and valence quark contribution (blue, dotted curve) and of the new GSF (green, dashed curve).

The corresponding, normalized, new GSF is plotted as the green, dashed curve in Fig. 2. It is the largest gluon Sivvers function compatible with PHENIX data: actually, it is strongly artificially enhanced by the extreme (or rather, unrealistic) assumption about the opposite contribution from saturated (and all summing up) sea-quark Sivvers functions. Nevertheless, it still indicates a rather modest GSF in the important small x region. Notice that the opposite choice, a negative contribution from the sea-quark Sivvers distributions balancing a positive one from the GSF, would lead to an even smaller GSF.

Finally, we notice that, concerning the BSR, within the x range covered by the data, the (over)maximized sea-quark Sivvers distributions give a positive contribution which strongly suppresses the negative contribution of the GSF, so that in this scenario the BSR is satisfied within a 10% level accuracy.

III. COMMENTS AND CONCLUSIONS

Unintegrated, TMD parton distribution and fragmentation functions may help in explaining several puzzling measurements for spin observables in inclusive particle production at high energy and moderately large p_T (for hadronic collisions) or Q^2 (for SIDIS). In particular, the quark Sivvers distributions and the Collins fragmentation functions have recently raised a lot of interest. Azimuthal and single spin asymmetries measured by several experimental collaborations unambiguously indicate that these effects are sizeable, at least in some kinematical regions.

A combined analysis of inclusive pion production in hadronic collisions and in SIDIS has given useful information on the quark Sivvers distributions in the valence region. On the contrary, sea-quark and gluon Sivvers functions are largely unknown. In this paper, we have performed a phenomenological analysis of the available data for the SSA $A_N(p^\uparrow p \rightarrow \pi X)$, in particular regarding the recent PHENIX data obtained at mid-rapidity and moderate p_T values; this SSA is dominated by the Sivvers effect and explores the low x region, resulting very sensitive to gluon contributions. To this end, we have adopted the theoretical approach of Refs. [1, 2, 3].

A first result of our analysis is that all available data, including the PHENIX ones, are compatible with valence-like quark Sivvers distributions and vanishing sea-quark and gluon contributions. The parameterizations required to reasonably reproduce the observed asymmetries also fulfill, within a 10% accuracy, the Burkardt sum rule. Since these parameterizations have a valence-like nature, the sum rule can be checked both over the fully integrated x range and in the limited x range effectively covered by data, with the same conclusions.

The main issue of this paper was, however, that of discussing to what extent the available data can bind the magnitude of the gluon Sivvers function, in particular in the small x region, where gluons play an important role. Our analysis shows that the PHENIX data on A_N are presently the only ones that allow to reach quantitative conclusions on the magnitude of the GSF. The weakest upper bound (that is, the largest GSF), is obtained by balancing the gluon contribution to A_N with (over)maximized sea-quark contributions, opposite in sign. The resulting GSF is represented by the green, dashed line of Fig. 2. This scenario corresponds to a SSA lying approximately within one-sigma deviation below the data, as shown by the thick, red, solid line of Fig. 1. Even in this extreme case, the bound obtained significantly reduces the magnitudes available from the simple positivity bound implicit in the definition of the Sivvers function, at least in the x region where gluon contributions are relevant.

Some comments are still in order:

- 1) Following Refs. [1, 2, 3], we have assumed for the TMD functions a simple factorized expression, with a Gaussian $k_\perp$ dependence. The same $\langle k_\perp^2 \rangle \simeq 0.64$ (GeV/c)² has been used for all partons (quarks and gluons). It has been suggested that gluons may have a larger $\langle k_\perp^2 \rangle$ than quarks [37]. We have found that, using smaller values of $\langle k_\perp^2 \rangle_q$

and, *e.g.*, $\langle k_{\perp}^2 \rangle_g \simeq 2 \langle k_{\perp}^2 \rangle_q$ would lead to a more stringent bound on the GSF.

2) In our calculations we have used the MRST01-LO set for distribution functions and the KKP-LO set of fragmentation functions. Since low- x values are relevant in the kinematical configuration of RHIC experiments, one may wonder whether the choice of unpolarized PDF's and FF's affects the bounds. We have checked that there is indeed some residual dependence of this type. As an example, using the CTEQ6 PDF's [38], which have a larger gluon component than the MRST01 set, would lead to a slightly more stringent bound. On the contrary, the use of the Kretzer FF set [39], which has a smaller gluon fragmentation component, would lead to a slightly less stringent bound. These small changes are well within the overall uncertainty of our results and do not alter our main conclusions.

Future measurements covering the negative x_F range, but at smaller c.m. energies, *e.g.* at RHIC with $\sqrt{s} = 63$ GeV or at the proposed PAX experiment at GSI, with $\sqrt{s} = 14$ GeV, will certainly help in clarifying the size and relevance of the GSF. SSA measurements for inclusive photon production might also give useful information. A detailed analysis of the phenomenological interest of these processes is in progress [40].

Acknowledgments

We acknowledge the support of the European Community–Research Infrastructure Activity under the FP6 “Structuring the European Research Area” programme (HadronPhysics, contract number RII3-CT-2004-506078); M.A. acknowledges partial support by MIUR under contract 2004021808.009.

Note added in proof

Just after submission of our paper an interesting work appeared [41] in which similar conclusions about the smallness of the gluon Sivers function are obtained from the smallness of the Sivers effect in SIDIS off a deuteron target.

-
- [1] U. D'Alesio and F. Murgia, Phys. Rev. **D70** (2004) 074009 [arXiv:hep-ph/0408092].
 - [2] M. Anselmino, M. Boglione, U. D'Alesio, E. Leader and F. Murgia, Phys. Rev. **D71** (2005) 014002 [arXiv:hep-ph/0408356].
 - [3] M. Anselmino, M. Boglione, U. D'Alesio, E. Leader, S. Melis and F. Murgia, Phys. Rev. **D73** (2006) 014020 [arXiv:hep-ph/0509035].
 - [4] M. Anselmino, M. Boglione, U. D'Alesio, E. Leader and F. Murgia, Phys. Rev. **D70** (2004) 074025 [arXiv:hep-ph/0407100].
 - [5] D.W. Sivers, Phys. Rev. **D41** (1990) 83; **D43** (1991) 261.
 - [6] J.C. Collins, Nucl. Phys. **B396** (1993) 161 [arXiv:hep-ph/9208213].
 - [7] J.C. Collins and D.E. Soper, Nucl. Phys. **B193** (1981) 381 [Erratum-ibid. **B213** (1983) 545].
 - [8] X.D. Ji, J.P. Ma and F. Yuan, Phys. Rev. **D71** (2005) 034005 [arXiv:hep-ph/0404183]; Phys. Lett. **B597** (2004) 299 [arXiv:hep-ph/0405085].
 - [9] J.C. Collins and A. Metz, Phys. Rev. Lett. **93** (2004) 252001 [arXiv:hep-ph/0408249].
 - [10] S.J. Brodsky, D.S. Hwang and I. Schmidt, Phys. Lett. **B530** (2002) 99 [arXiv:hep-ph/0201296]; Nucl. Phys. **B642** (2002) 344 [arXiv:hep-ph/0206259].
 - [11] J.C. Collins, Phys. Lett. **B536** (2002) 43 [arXiv:hep-ph/0204004].
 - [12] A.V. Belitsky, X. Ji and F. Yuan, Nucl. Phys. **B656** (2003) 165 [arXiv:hep-ph/0208038].
 - [13] X.D. Ji and F. Yuan, Phys. Lett. **B543** (2002) 66 [arXiv:hep-ph/0206057].
 - [14] C.J. Bomhof, P.J. Mulders and F. Pijlman, Phys. Lett. **B596** (2004) 277 [arXiv:hep-ph/0406099].
 - [15] A. Bacchetta, C.J. Bomhof, P.J. Mulders and F. Pijlman, Phys. Rev. **D72** (2005) 034030 [arXiv:hep-ph/0505268].
 - [16] A. Airapetian *et al.* (HERMES Collaboration), Phys. Rev. Lett. **94** (2005) 012002 [arXiv:hep-ex/0408013]; M. Dieffenhafer (for the HERMES Collaboration), AIP Conf. Proc. **792** (2005) 933 [arXiv:hep-ex/0507013].
 - [17] V.Y. Alexakhin *et al.* (COMPASS Collaboration), Phys. Rev. Lett. **94** (2005) 202002 [arXiv:hep-ex/0503002].
 - [18] M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, F. Murgia and A. Prokudin, Phys. Rev. **D71** (2005) 074006 [arXiv:hep-ph/0501196]; **D72** (2005) 094007 [arXiv:hep-ph/0507181].
 - [19] W. Vogelsang and F. Yuan, Phys. Rev. **D72** (2005) 054028 [arXiv:hep-ph/0507266].
 - [20] A.V. Efremov, K. Goeke, S. Menzel, A. Metz and P. Schweitzer, Phys. Lett. **B612** (2005) 233 [arXiv:hep-ph/0412353].
 - [21] J.C. Collins, A.V. Efremov, K. Goeke, S. Menzel, A. Metz and P. Schweitzer, Phys. Rev. **D73** (2006) 014021 [arXiv:hep-ph/0509076].
 - [22] M. Anselmino *et al.*, arXiv:hep-ph/0511017.
 - [23] P.J. Mulders and J. Rodrigues, Phys. Rev. **D63** (2001) 094021 [arXiv:hep-ph/0009343].
 - [24] D. Boer and W. Vogelsang, Phys. Rev. **D69** (2004) 094025 [arXiv:hep-ph/0312320].
 - [25] I. Schmidt, J. Soffer and J.J. Yang, Phys. Lett. **B612** (2005) 258 [arXiv:hep-ph/0503127].
 - [26] D.L. Adams *et al.* (FNAL-E704 Collaboration), Phys. Lett. **B264** (1991) 462; A. Bravar *et al.* (FNAL-E704 Collaboration), Phys. Rev. Lett. **77** (1996) 2626.




FIG. 1: The SSA A_N , computed according to Eqs. (1) and (3)-(5) of the text, and compared with PHENIX data [28], with different choices for the gluon and sea-quark Sivers functions. The thin, red, solid line shows the results given by the valence u and d Sivers functions alone. The cyan, dot-dashed curve shows the contribution of the maximized GSF alone, $\Delta^N \hat{f}_{g/p\uparrow} = -2\hat{f}_{g/p}$. The thick, red, solid curve is obtained by requesting results within one standard deviation from data in two different ways: no sea-quark contribution and maximized sea-quark contribution. In the latter case the blue, dotted curve shows the contribution of the sea (maximized) + valence quarks, while the green, dashed curve shows the contribution of the GSF. The valence quark Sivers distributions are taken from Ref. [1], while the unpolarized PDF's and FF's from Refs. [35] and [36] respectively. Further detail can be found in the text.

- [26] J. Adams *et al.* (STAR Collaboration), Phys. Rev. Lett. **92** (2004) 171801 [arXiv:hep-ex/0310058].
- [27] D.L. Adams *et al.* (FNAL-E704 Collaboration), Phys. Rev. **D53** (1996) 4747.
- [28] S.S. Adler *et al.* (PHENIX Collaboration), Phys. Rev. Lett. **95** (2005) 202001 [arXiv:hep-ex/0507073].
- [29] C.A. Gagliardi (for the STAR Collaboration), arXiv:hep-ex/0607003.
- [30] F. Videbaek (for the BRAHMS Collaboration), arXiv:nucl-ex/0601008.
- [31] V. Barone *et al.* (PAX Collaboration), arXiv:hep-ex/0505054.
- [32] U. D'Alesio and F. Murgia, unpublished.
- [33] M. Burkardt, Phys. Rev. **D69** (2004) 057501 [arXiv:hep-ph/0311013]; **D69** (2004) 091501 [arXiv:hep-ph/0402014].
- [34] K. Goeke, S. Meissner, A. Metz and M. Schlegel, Phys. Lett. **B637** (2006) 241 [arXiv:hep-ph/0601133].
- [35] A.D. Martin, R.G. Roberts, W.J. Stirling and R.S. Thorne, Phys. Lett. **B531** (2002) 216 [arXiv:hep-ph/0201127].
- [36] B.A. Kniehl, G. Kramer and B. Potter, Nucl. Phys. **B582** (2000) 514 [arXiv:hep-ph/0010289].
- [37] W. Vogelsang, private communication.
- [38] J. Pumplin, D.R. Stump, J. Huston, H.L. Lai, P. Nadolsky and W.K. Tung, JHEP **0207** (2002) 012 [arXiv:hep-ph/0201195].
- [39] S. Kretzer, Phys. Rev. **D62** (2000) 054001 [arXiv:hep-ph/0003177].
- [40] U. D'Alesio, S. Melis and F. Murgia, in preparation.
- [41] S.J. Brodsky and S. Gardner, arXiv:hep-ph/0608219.




FIG. 2: The value of the normalized GSF, $|\Delta^N f_{g/p^\uparrow}(x)|/2f_{g/p}(x)$, as obtained from fitting, within one standard deviation, the PHENIX data on A_N . The red, solid curve corresponds to the case of vanishing sea-quark contribution; the green, dashed curve shows the GSF corresponding to the extreme scenario in which the sea-quark contributions are all maximized and sum up to balance the negative GSF contribution. Further detail can be found in the text.